arXiv:1104.2835v2 [math.AC] 17 Apr 2011

On glued semigroups

J. I. García-García*
M.A. Moreno-Frías†
A. Vigneron-Tenorio‡

January 19, 2013

Abstract

This paper presents some results about the combinatorial properties of the glued semigroups and gives a combinatorial characterization of them. Some classical results for affine glued semigroups are generalized to finitely generated commutative reduced cancellative and with zero element glued semigroups. In addition, some algorithms to construct this kind of semigroups with different properties are given.

Keywords: Semigroup, semigroup ideal, gluing of semigroups, simplicial complex, toric ideal.

MSC-class: 13F20 (Primary) 05E40, 13F55 (Secondary).

Introduction

Let $S=\langle n_1,\dots,n_l\rangle$ be a finitely generated commutative semigroup with zero element such that it is reduced (i.e. $S\cap(-S)=(0)$). We suppose that S is cancellative, that is to say, if $m+n=m+n'$, with $m,n,n'\in S$, then $n=n'$. With these conditions, we may assume that S is a subsemigroup of a non necessarily torsion-free group. If S is torsion-free, then S is an affine semigroup.

From now on, we assume that all the semigroups appearing in this work are finitely generated, commutative and reduced, thus in the sequel we omit these adjectives.

Let $\Bbbk$ be a field and $\Bbbk[X_1,\dots,X_l]$ the polynomial ring in l indeterminates. This polynomial ring is obviously an $S-$graded ring (by assigning the S-degree n_i to the indeterminate X_i, the S-degree of $X^\alpha=X_1^{\alpha_1}\cdots X_l^{\alpha_l}$ is $\sum_{i=1}^l\alpha_i n_i\in S$).

---

*Departamento de Matemáticas, Universidad de Cádiz, E-11510 Puerto Real (Cádiz, Spain). E-mail: ignacio.garcia@uca.es. Partially supported by MTM2007-62346 and Junta de Andalucía group FQM-366.

†Departamento de Matemáticas, Universidad de Cádiz, E-11510 Puerto Real (Cádiz, Spain). E-mail: mariangeles.moreno@uca.es. Partially supported by MTM2008-06201-C02-02 and Junta de Andalucía group FQM-298.

‡Departamento de Matemáticas, Universidad de Cádiz, E-11405 Jerez de la Frontera (Cádiz, Spain). E-mail: alberto.vigneron@uca.es. Partially supported by MTM2007-64704 and Junta de Andalucía group FQM-366.

It is well known that the ideal (denoted by I_S) generated by

$$\left\{ X^\alpha - X^\beta | \sum_{i=1}^{l} \alpha_i n_i = \sum_{i=1}^{l} \beta_i n_i \right\} \subset \Bbbk[X_1, \dots, X_l]$$

is an $S-$homogeneous binomial ideal called *semigroup ideal* (see [6] for details). If S is torsion-free, the ideal obtained defines a toric variety (see [12] and the references therein). By Nakayama's lemma, all minimal generating sets of I_S have same cardinality and the $S-$degrees of its elements are determinated.

In [1], [4] and [7] the authors study the minimal generating sets of semigroup ideals by means of the homology of different simplicial complexes (with isomorphic homologies) associated to the semigroup. For any $m \in S$, set

$$C_m = \{X^\alpha = X_1^{\alpha_1} \cdots X_l^{\alpha_l} \mid \sum_{i=1}^{l} \alpha_i n_i = m\}, \tag{1}$$

we consider the abstract simplicial complex (used in [4] and [7]) on the vertex set C_m,

$$\nabla_m = \{F \subseteq C_m \mid \gcd(F) \neq 1\}, \tag{2}$$

where $\gcd(F)$ is the *greatest common divisor* of the monomials in F.

The main aim of this work is to study the semigroups which result from the gluing of other two. This concept was introduced by Rosales in [10], and it is closely related to the ideals that are complete intersections (see [13] and the references therein).

A semigroup S minimally generated by $A_1 \sqcup A_2$ (with $A_1 = \{n_1, \dots, n_r\}$ and $A_2 = \{n_{r+1}, \dots, n_l\}$) is the gluing of $S_1 = \langle A_1 \rangle$ and $S_2 = \langle A_2 \rangle$, if there exists a set of generators, ρ, of I_S of the form

$$\rho = \rho_1 \cup \rho_2 \cup \{X^\gamma - X^{\gamma'}\},$$

where ρ_1, ρ_2 are generating sets of I_{S_1} and I_{S_2} respectively, and $X^\gamma - X^{\gamma'} \in I_S$ such that the support of γ (supp (γ)) is included in $\{1, \dots, r\}$ and supp $(\gamma') \subset \{r+1, \dots, l\}$. Equivalently, S is the gluing of S_1 and S_2 if $I_S = I_{S_1} + I_{S_2} + \langle X^\gamma - X^{\gamma'} \rangle$. We call *glued semigroups* to this kind of semigroups.

In Section 1, we define the required mathematical elements in order to generalize to non torsion-free semigroups a classical result concerning affine semigroups (Proposition 2).

In Section 2, we examine the non-connected simplicial complexes ∇_m associated to the glued semigroups. By understanding the vertices of the connected components of these complexes, we give a combinatorial characterization of the glued semigroups as well as their glued degrees (Theorem 6). Besides, in Corollary 7 we deduce the conditions under which the ideal of a glued semigroup is uniquely generated. Despite the fact that Theorem 6 and Corollary 7 provide the basis to implement algorithms, they may be, however, no efficient. In this sense, the goal of this section is to provide further knowledge about (glued) semigroups employing combinatorial theory no matter the efficiency of the obtained algorithms.

We devote the last part of this work, Section 3, to construct glued semigroups (Corollary 10), complete intersection glued semigroups and affine glued semigroups (Subsection 3.1). We create the affine glued semigroups by solving an integer programming problem.

1 Preliminaries and generalizations about glued semigroups

In this section we summarize some notations and definitions, and give a generalization to non-torsion free semigroups of [10, Theorem 1.4].

We say that a binomial in I_S is *indispensable* if it is in all the system of generators of I_S (up to a scalar multiple). This kind of binomials were introduced in [9]. This notion comes from the Algebraic Statistics. In [8] the authors characterize the indispensable binomials by using the simplicial complexes ∇_m. Note that if I_S is generated by its indispensable binomials, I_S is uniquely generated (up to a scalar multiple).

Being the notation set as in the introduction, we associate a lattice to the semigroup S: $\ker S \subset \mathbb{Z}^l$, $\alpha = (\alpha_1, \ldots, \alpha_l) \in \ker S$ if $\sum_{i=1}^{l} \alpha_i n_i = 0$. The property *"S is reduced"* is equivalent to $\ker S \cap \mathbb{N}^l = (0)$. Given a system of binomial generators of I_S, $\ker S$ is generated by a set whose elements are $\alpha - \beta$ with $X^\alpha - X^\beta$ being in the system of binomial generators.

We call $\mathcal{M}(I_S)$ to a minimal generating set of I_S, and $\mathcal{M}(I_S)_m \subset \mathcal{M}(I_S)$ to the set of their elements whose $S-$degree are equal to $m \in S$. *Betti*(S) is the set of the $S-$degrees of the elements in $\mathcal{M}(I_S)$.

S is called a *complete intersection* semigroup if I_S is minimally generated by $\text{rank}(\ker S)$ elements.

Let $\mathcal{C}(\nabla_m)$ be the number of connected components of a non-connected ∇_m, this means that the cardinality of $\mathcal{M}(I_S)_m$ is $\mathcal{C}(\nabla_m) - 1$ (see Remark 2.6 in [1] and Theorem 3 and Corollary 4 in [7]). Note that the complexes associated to the elements in *Betti*(S) are non-connected. The relation between $\mathcal{M}(I_S)$ and *Betti*(S) is studied next.

Construction 1. ([4, Proposition 1]). *For each $m \in$ Betti(S), one can construct $\mathcal{M}(I_S)_m$ by taking $\mathcal{C}(\nabla_m) - 1$ binomials whose monomials are in different connected components of ∇_m and satisfying that two different binomials have not their corresponding monomials in the same components. This let us construct a minimal generating set of I_S in a combinatorial way.*

Now, we are going to introduce the notations that we use to work with glued semigroups.

Let S be minimally[1] generated by $A_1 \sqcup A_2$ with $A_1 = \{a_1, \ldots, a_r\}$ and $A_2 = \{b_1, \ldots, b_t\}$. From now on, we identify the sets A_1 and A_2 with the matrixes

$$\begin{pmatrix} a_1 \\ \vdots \\ a_r \end{pmatrix} \text{ and } \begin{pmatrix} b_1 \\ \vdots \\ b_t \end{pmatrix}.$$

We denote by $\Bbbk[A_1]$ and $\Bbbk[A_2]$ to the polinomial rings $\Bbbk[X_1, \ldots, X_r]$ and $\Bbbk[Y_1, \ldots, Y_t]$, respectively. We call *pure monomials* to the monomials with indeterminates only in $X_1, \ldots, X_r$ or $Y_1, \ldots, Y_t$. Conversely, we call *mixed monomials* to the monomials with indeterminates in Xs and Ys.

---

[1] We consider a minimal generator set of S because, in the other case, S is trivially gluing of the semigroup generated by one of its non minimal generators and the semigroup generated by the others.

Given S, the gluing of $S_1 = \langle A_1 \rangle$ and $S_2 = \langle A_2 \rangle$, we say that $X^{\gamma_X} - Y^{\gamma_Y} \in I_S$ is a *glued binomial* if $\mathcal{M}(I_{S_1}) \cup \mathcal{M}(I_{S_2}) \cup \{X^{\gamma_X} - Y^{\gamma_Y}\}$ is a generating set of I_S. The element $d = S$-*degree*$(X^{\gamma_X}) \in S$ is called a *glued degree.*

In this way, it is clear that if S is a glued semigroup, the lattice $\ker S$ has a basis such as

$$\{L_1, L_2, (\gamma_X, -\gamma_Y)\} \subset \mathbb{Z}^{r+t}, \tag{3}$$

where the supports of the elements in L_1 are in $\{1, \ldots, r\}$, the supports of the elements in L_2 are in $\{r+1, \ldots, r+t\}$, $\ker S_i = L_i\mathbb{Z}\,(i = 1, 2)$ by considering only the coordinates in $\{1, \ldots, r\}$ or $\{r+1, \ldots, r+t\}$ of L_i, and $(\gamma_X, \gamma_Y) \in \mathbb{N}^{r+t}$. Moreover, since S is reduced, one has that $L_1\mathbb{Z} \cap \mathbb{N}^{r+t} = L_2\mathbb{Z} \cap \mathbb{N}^{r+t} = (0)$. We will denote by $\{\rho_{1i}\}_i$ to the elements in L_1 and by $\{\rho_{2i}\}_i$ to the elements in L_2.

The following Proposition generalizes [10, Theorem 1.4] to non-torsion free semigroups.

Proposition 2. *S is the gluing of S_1 and S_2 if and only if there exists $d \in (S_1 \cap S_2) \setminus \{0\}$ such that $G(S_1) \cap G(S_2) = d\mathbb{Z}$, where $G(S_1)$, $G(S_2)$ and $d\mathbb{Z}$ are the associated commutative groups of S_1, S_2 and $\{d\}$.*

Proof. Let's assume that S is the gluing of S_1 and S_2. In this case, $\ker S$ is generated by the set (3). Since $(\gamma_X, -\gamma_Y) \in \ker S$, $d = A_1\gamma_X = A_2\gamma_Y \in S$ and $d \in S_1 \cap S_2 \subset G(S_1) \cap G(S_2)$.

Let d' be in $G(S_1) \cap G(S_2)$, then $\exists(\delta_1, \delta_2) \in \mathbb{Z}^r \times \mathbb{Z}^t$ such that $d' = A_1\delta_1 = A_2\delta_2$. Therefore $(\delta_1, -\delta_2) \in \ker S$ because $(A_1|A_2)(\delta_1, -\delta_2) = 0$. By taking this into consideration, we find that there exist $\lambda, \lambda_i^{\rho_1}, \lambda_i^{\rho_2} \in \mathbb{Z}$ satisfying

$$\begin{cases} (\delta_1, 0) &= \quad \sum_i \lambda_i^{\rho_1} \rho_{1i} + \lambda(\gamma_X, 0) \\ \\ (0, \delta_2) &= -\sum_i \lambda_i^{\rho_2} \rho_{2i} + \lambda(0, \gamma_Y), \end{cases}$$

and $d' = A_1\delta_1 = \sum_i \lambda_i^{\rho_1}(A_1|0)\rho_{1i} + \lambda A_1\gamma_X = \lambda d$. Therefore, we conclude that $G(S_1) \cap G(S_2) = d\mathbb{Z}$ with $d \in S_1 \cap S_2$.

Conversely, we suppose that there exists $d \in (S_1 \cap S_2) \setminus \{0\}$ such that $G(S_1) \cap G(S_2) = d\mathbb{Z}$. Assuming this, we will prove that $I_S = I_{S_1} + I_{S_2} + \langle X^{\gamma_X} - Y^{\gamma_Y} \rangle$. Trivially, $I_{S_1} + I_{S_2} + \langle X^{\gamma_X} - Y^{\gamma_Y} \rangle \subset I_S$.

Let $X^\alpha Y^\beta - X^\gamma Y^\delta$ be a binomial in I_S. Its $S-$degree is $A_1\alpha + A_2\beta = A_1\gamma + A_2\delta$. Since $A_1(\alpha - \gamma) = A_2(\beta - \delta) \in G(S_1) \cap G(S_2) = d\mathbb{Z}$, there exists $\lambda \in \mathbb{Z}$ such that $A_1\alpha = A_1\gamma + \lambda d$ and $A_2\delta = A_2\beta + \lambda d$.

- If $\lambda = 0$,

$$X^\alpha Y^\beta - X^\gamma Y^\delta = X^\alpha Y^\beta - X^\gamma Y^\beta + X^\gamma Y^\beta - X^\gamma Y^\delta =$$
$$= Y^\beta(X^\alpha - X^\gamma) + X^\gamma(Y^\beta - Y^\delta) \in I_{S_1} + I_{S_2}.$$

- If $\lambda > 0$,

$$X^\alpha Y^\beta - X^\gamma Y^\delta =$$
$$= X^\alpha Y^\beta - X^\gamma X^{\lambda\gamma_X} Y^\beta + X^\gamma X^{\lambda\gamma_X} Y^\beta - X^\gamma X^{\lambda\gamma_Y} Y^\beta + X^\gamma X^{\lambda\gamma_Y} Y^\beta - X^\gamma Y^\delta =$$
$$= Y^\beta(X^\alpha - X^\gamma X^{\lambda\gamma_X}) + X^\gamma Y^\beta(X^{\lambda\gamma_X} - Y^{\lambda\gamma_Y}) + X^\gamma(Y^{\lambda\gamma_Y} Y^\beta - Y^\delta).$$

  This last polynomial is in $I_{S_1} + I_{S_2} + \langle X^{\gamma_X} - Y^{\gamma_Y} \rangle$ because of

$$X^{\lambda\gamma_X} - Y^{\lambda\gamma_Y} = (X^{\gamma_X} - Y^{\gamma_Y}) \left(\sum_{i=1}^{\lambda-1} X^{(\lambda-i)\gamma_X} Y^{i\gamma_Y} \right).$$

- The case $\lambda < 0$ can be solved likewise.

Therefore, we conclude that $I_S = I_{S_1} + I_{S_2} + \langle X^{\gamma_X} - Y^{\gamma_Y}\rangle$.

$\square$

It follows that, given the partition of the system of generators of S, the glued degree is unique.

2 Glued semigroups and combinatorics

In this section, we approach the study of simplicial complexes ∇_m associated with glued semigroups. We characterize the glued semigroups by means of the non-connected simplicial complexes.

For any $m \in S$, we redefine C_m from (1), as

$$C_m = \{X^\alpha Y^\beta = X_1^{\alpha_1} \cdots X_r^{\alpha_r} Y_1^{\beta_1} \cdots Y_t^{\beta_t} \mid \sum_{i=1}^{r} \alpha_i a_i + \sum_{i=1}^{t} \beta_i b_i = m\}$$

and consider the vertex sets and the simplicial complexes

$$C_m^{A_1} = \{X_1^{\alpha_1} \cdots X_r^{\alpha_r} \mid \sum_{i=1}^{r} \alpha_i a_i = m\},\ \nabla_m^{A_1} = \{F \subseteq C_m^{A_1} \mid \gcd(F) \neq 1\},$$

$$C_m^{A_2} = \{Y_1^{\beta_1} \cdots Y_t^{\beta_t} \mid \sum_{i=1}^{t} \beta_i b_i = m\},\ \nabla_m^{A_2} = \{F \subseteq C_m^{A_2} \mid \gcd(F) \neq 1\},$$

where $A_1 = \{a_1, \ldots, a_r\}$ and $A_2 = \{b_1, \ldots, b_t\}$ as in Section 1.

Trivially, the relations between $\nabla_m^{A_1}, \nabla_m^{A_2}$ and ∇_m are

$$\nabla_m^{A_1} = \{F \in \nabla_m | F \subset C_m^{A_1}\},\ \nabla_m^{A_2} = \{F \in \nabla_m | F \subset C_m^{A_2}\}. \tag{4}$$

The following result shows a relevant property of the simplicial complexes associated to glued semigroups.

Lemma 3. *Let S be the gluing of S_1 and S_2, and $m \in Betti(S)$. Then all the connected components of ∇_m have at least a pure monomial. In addition, all mixed monomials of ∇_m are in the same connected component.*

Proof. Supposed that there exists C, a connected component of ∇_m only with mixed monomials. In this case, in any generating set of I_S there is, at least, a binomial with a mixed monomial (by Construction 1). But there is not this kind of binomial in $\mathcal{M}(I_{S_1}) \cup \mathcal{M}(I_{S_2}) \cup \{X^{\gamma_X} - Y^{\gamma_Y}\}$, with $X^{\gamma_X} - Y^{\gamma_Y}$ a glued binomial. This is not possible because S is the gluing of S_1 and S_2.

Since S is a glued semigroup, $\ker S$ has a system of generators as the introduced in (3). Let $X^\alpha Y^\beta, X^\gamma Y^\delta \in C_m$ be two monomials such that $\gcd(X^\alpha Y^\beta, X^\gamma Y^\delta) = 1$. In this case, $(\alpha, \beta) - (\gamma, \delta) \in \ker S$, and $\exists \lambda, \lambda_i^{\rho_1}, \lambda_i^{\rho_2} \in \mathbb{Z}$ satisfying:

$$\begin{cases} (\alpha - \gamma, 0) &= \sum_i \lambda_i^{\rho_1} \rho_{1i} + \lambda(\gamma_X, 0) \\ (0, \beta - \delta) &= \sum_i \lambda_i^{\rho_2} \rho_{2i} - \lambda(0, \gamma_Y) \end{cases}$$

- If $\lambda = 0$, $\alpha - \gamma \in \ker S_1$ and $\beta - \delta \in \ker S_2$. Then $A_1\alpha = A_1\gamma$, $A_2\beta = A_2\delta$ and $X^\alpha Y^\delta \in C_m$.
- If $\lambda > 0$, $(\alpha, 0) = \sum_i \lambda_i^{\rho_1} \rho_{1i} + \lambda(\gamma_X, 0) + (\gamma, 0)$ and
$$A_1\alpha = \sum_i \lambda_i^{\rho_1}(A_1|0)\rho_{1i} + \lambda A_1\gamma_X + A_1\gamma = \lambda d + A_1\gamma.$$
  Then $X^{\lambda\gamma_X} X^\gamma Y^\beta \in C_m$.
- The case $\lambda < 0$ can be solved likewise.

In any case, $X^\alpha Y^\beta$ and $X^\gamma Y^\delta$ are in the same connected component of ∇_m. $\square$

The following Lemma describes the simplicial complexes that correspond to the $S-$degrees that are multiples of the glued degree.

Lemma 4. *Let S be the gluing of S_1 and S_2, $d \in S$ the glued degree, and $d' \in S$. Then $C_{d'}^{A_1} \neq \emptyset \neq C_{d'}^{A_2}$ if and only if $d' \in (d\mathbb{N}) \setminus \{0\}$. Furthermore, the simplicial complex $\nabla_{d'}$ has at least one connected component with elements in $C_{d'}^{A_1}$ and $C_{d'}^{A_2}$.*

Proof. Supposed that there exist $X^\alpha, Y^\beta \in C_{d'}$, then $d' = \sum_{i=1}^r \alpha_i a_i = \sum_{i=1}^t \beta_i b_i \in S_1 \cap S_2 \subset G(S_1) \cap G(S_2) = d\mathbb{Z}$. Trivially, $d' \in d\mathbb{N}$.

Conversely, $d' = jd$, with $j \in \mathbb{N}$. Let $X^{\gamma_X} - Y^{\gamma_Y} \in I_S$ be a glued binomial. It is easy to see that $X^{j\gamma_X}, Y^{j\gamma_Y} \in C_{d'}$. Trivially, $\{X^{j\gamma_X}, X^{(j-1)\gamma_X} Y^{\gamma_Y}\}$ and $\{X^{(j-1)\gamma_X} Y^{\gamma_Y}, Y^{j\gamma_Y}\}$ are in $\nabla_{d'}$. $\square$

The following Lemma is a combinatorial version of [5, Lemma 9] and it is a necessary condition for our combinatorial characterization theorem (Theorem 6).

Lemma 5. *Let S be the gluing of S_1 and S_2, and $d \in S$ the glued degree. Then the elements in C_d are pure monomials and $d \in Betti(S)$.*

Proof. The reader can check that $m' \preceq_S m$ if $m - m' \in S$, is a well defined partial order on S.

Let's assume that there exists a mixed monomial $T \in C_d$. By Lemma 3, there exists a pure monomial in C_d, for example Y^b, such that $\{T, Y^b\} \in \nabla_d$ (this proof is analogous if we consider X^a such that $\{T, X^a\} \in \nabla_d$). Thus, if we get $T_1 = \gcd(T, Y^b)^{-1} T$ and $Y^{b_1} = \gcd(T, Y^b)^{-1} Y^b$, we conclude that both monomials are in $C_{d'}$, where d' is equal to d minus the $S-$degree of $\gcd(T, Y^b)$.

If $C_{d'}^{A_1} \neq \emptyset$, by Lemma 4, $d' \in d\mathbb{N}$. But this is not possible due to the fact that $d' \prec_S d$. Consequently, one can consider that T_1 is a mixed monomial and $C_{d'}^{A_1} = \emptyset$, but $C_{d'}^{A_2}$ is not empty.

If there exists a pure monomial in $C_{d'}^{A_2}$ connected to a mixed monomial in $C_{d'}$, the above process can be repeated until $T_2, Y^{b_2} \in C_{d''}$ are obtained, with T_2 a mixed monomial. This process is finite by degree reasons.

So, after some steps, one find a $d^{(i)} \in S$ such that $\nabla_{d^{(i)}}$ is not connected (i.e. $d^{(i)} \in Betti(S)$) and it has a connected component whose vertices are only mixed monomials. This contradicts Lemma 3. $\square$

After examining the structure of the simplicial complexes associated to the glued semigroups, we enunciate a combinatorial characterization theorem by means of the non-connected simplicial complexes ∇_m. This result helps to understand the nature of glued semigroups and increase our knowledge on them.

Theorem 6. *S is the gluing of S_1 and S_2 if and only if:*

1. *For all $d' \in Betti(S)$, any connected component of $\nabla_{d'}$ has at least a pure monomial.*
2. *There exists an unique $d \in Betti(S)$ such that $C_d^{A_1} \neq \emptyset \neq C_d^{A_2}$ and the elements in C_d are pure monomials.*
3. *For all $d' \in Betti(S) \setminus \{d\}$ with $C_{d'}^{A_1} \neq \emptyset \neq C_{d'}^{A_2}$, we have $d' \in d\mathbb{N}$.*

Besides, the above $d \in Betti(S)$ is the glued degree.

Proof. If S is the gluing of S_1 and S_2, we obtain immediately the theorem from Lemmas 3, 4 and 5.

Conversely, given $d' \in Betti(S) \setminus \{d\}$, by the hypothesis 1 and 3, we can construct the sets, $\mathcal{M}(I_{S_1})_{d'}$ and $\mathcal{M}(I_{S_2})_{d'}$, in a similar way to Construction 1, but only taking binomials whose monomials are in $C_{d'}^{A_1}$ or $C_{d'}^{A_2}$.

Analogously, if we consider $d \in Betti(S)$, we construct the set $\mathcal{M}(I_S)_d$ with $\mathcal{C}(\nabla_d) - 1$ binomials as the union of:

- $\mathcal{C}(\nabla_d^{A_1}) - 1$ binomials in $\mathcal{M}(I_{S_1})_d$.
- $\mathcal{C}(\nabla_d^{A_2}) - 1$ binomials in $\mathcal{M}(I_{S_2})_d$.
- A binomial $X^{\gamma_X} - Y^{\gamma_Y}$, with $X^{\gamma_X} \in C_d^{A_1}$ and $Y^{\gamma_Y} \in C_d^{A_2}$.

We conclude that $\bigsqcup_{m \in Betti(S)} \Big(\mathcal{M}(I_{S_1})_m \sqcup \mathcal{M}(I_{S_2})_m\Big) \sqcup \{X^{\gamma_X} - Y^{\gamma_Y}\}$ is a generating set of I_S. So S is the gluing of S_1 and S_2. □

From Theorem 6 we obtain an equivalent property to that in [5, Theorem 12] by using the *language* of monomials and binomials.

Corollary 7. *Let S be the gluing of S_1 and S_2, and $X^{\gamma_X} - Y^{\gamma_Y} \in I_S$ a glued binomial with $S-$degree d. Then, I_S is (minimally) generated by its indispensable binomials if and only if:*

- *I_{S_1} and I_{S_2} are (minimally) generated by their indispensable binomials.*
- *$X^{\gamma_X} - Y^{\gamma_Y}$ is an indispensable binomial of I_S.*
- *For all $d' \in Betti(S)$, the elements of $C_{d'}$ are pure monomials.*

Proof. Suppose that I_S is generated by its indispensable binomials. By [8, Corollary 6], $\forall m \in Betti(S)$, ∇_m has only two vertices. In particular, by Theorem 1, $\nabla_d = \{\{X^{\gamma_X}\}, \{Y^{\gamma_Y}\}\}$, and for $d' \in Betti(S) \setminus \{d\}$, $\nabla_{d'}$ is equal to $\nabla_{d'}^{A_1}$ or $\nabla_{d'}^{A_2}$ (by Lemma 1).

In any case, $X^{\gamma_X} - Y^{\gamma_Y} \in I_S$ is an indispensable binomial, and I_{S_1}, I_{S_2} are generated by their indispensable binomials.

Conversely, we suppose that I_S is not generated by its indispensable binomials. So, $\exists d' \in Betti(S) \setminus \{d\}$ such that $\nabla_{d'}$ has more than two vertices in at least two different connected components. Taking into account our hypothesis, there are not mixed monomials in $\nabla_{d'}$ and so:

- If $\nabla_{d'}$ is equal to $\nabla_{d'}^{A_1}$ (or $\nabla_{d'}^{A_2}$), then I_{S_1} (or I_{S_2}) is not generated by its indispensable binomials.
- In other case, $C_{d'}^{A_1} \neq \emptyset \neq C_{d'}^{A_2}$, by Lemma 4, $d' = jd$, with $j \in \mathbb{N}$, and so $X^{(j-1)\gamma_X} Y^{\gamma_Y} \in C_{d'}$, which contradicts our hypothesis.

Thus, we conclude that I_S is generated by its indispensable binomials. □

We illustrate the above results with the following example taken from [13].

Example 8. *Let $S \subset \mathbb{N}^2$ be the semigroup generated by the set*

$$\{(13,0),(5,8),(2,11),(0,13),(4,4),(6,6),(7,7),(9,9)\}.$$

In this case, Betti(S) is

$$\{(15,15),(14,14),(12,12),(18,18),(10,55),(15,24),(13,52),(13,13)\}.$$

Using the appropriated notation for the indeterminates in the polinomial ring $\Bbbk[x_1,\ldots,x_4,y_1,\ldots,y_4]$ (x_1,x_2,x_3 and x_4 for the first four generators of S and y_1,y_2,y_3,y_4 for the others), one can see that the simplicial complexes associated to the elements in Betti(S) are those that appear in Figure 1.


$C_{(15,15)} = \{y_1^2y_3, y_2y_4\}$
$\nabla_{(15,15)}$
Ys
$C_{(14,14)} = \{y_1^2y_2, y_3^2\}$
$\nabla_{(14,14)}$
Ys
$C_{(12,12)} = \{y_1^3, y_2^2\}$
$\nabla_{(12,12)}$
Ys
$C_{(10,55)} = \{x_1^2x_4, x_3^5\}$
$\nabla_{(10,55)}$
Xs
$C_{(18,18)} = \{y_1^2y_2, y_2^3, y_1y_3^2, y_4^2\}$
$\nabla_{(18,18)}$
Ys
$C_{(15,24)} = \{x_1x_2x_3, x_2^3, x_3y_1y_4, x_3y_2y_3\}$
$\nabla_{(15,24)}$
Ys
Xs
$C_{(13,52)} = \{x_2x_3^4, x_1x_4^4, x_4^3y_1y_4, x_4^3y_2y_3\}$
$\nabla_{(13,52)}$
Ys
Xs
$C_{(13,13)} = \{x_1x_4, y_1y_4, y_2y_3\}$
$\nabla_{(13,13)}$
Ys
Xs

Figure 1: Non-connected simplicial complexes associated to *Betti*(S).

From Figure 1, by using Theorem 6, one can concluded that S is the gluing of the semigroups $\langle(13,0),(5,8),(2,11),(0,13)\rangle$ and $\langle(4,4),(6,6),(7,7),(9,9)\rangle$, and the glued degree is $(13,13)$.

From Corollary 7, I_S is not generated by its indispensable binomials (I_S has only four indispensable binomials).

3 Generating glued semigroups

In this section, we give an algorithm with the aim of producing many examples of glued semigroups. Furthermore, we construct affine glued semigroups by means of solving an integer programming problem.

First of all, we consider two semigroups T_1 and T_2. Keeping the same notation we have followed throughout the whole article, let $A_1 = \{a_1, \ldots, a_r\}$ and $A_2 = \{b_1, \ldots, b_t\}$ be two minimal generator sets of the semigroups $T_1 = \langle A_1 \rangle$ and $T_2 = \langle A_2 \rangle$, and $L_j = \{\rho_{ji}\}_i$ be a basis of $\ker T_j$ with $j = 1, 2$.

Let γ_X and γ_Y be two nonzero elements in $\mathbb{N}^r$ and $\mathbb{N}^t$ respectively[2], and consider the integer matrix

$$A = \left(\begin{array}{c|c} L_1 & 0 \\ \hline 0 & L_2 \\ \hline \gamma_X & -\gamma_Y \end{array} \right). \tag{5}$$

Let S be a semigroup such that $\ker S$ is the lattice generated by the rows of the matrix A. This semigroup can be computed by using the Smith Normal Form (see [11, Chapter 2]). Denote by B_1, B_2 to two sets of cardinality r and t respectively, satisfying $S = \langle B_1, B_2 \rangle$ and $\ker(\langle B_1, B_2 \rangle)$ is generated by the rows of A.

The following Proposition shows that the semigroup S satisfies one of the conditions to be a glued semigroup.

Proposition 9. *The semigroup S verifies $G(\langle B_1 \rangle) \cap G(\langle B_2 \rangle) = (B_1\gamma_X)\mathbb{Z} = (B_2\gamma_Y)\mathbb{Z}$ with $d = B_1\gamma_X \in \langle B_1 \rangle \cap \langle B_2 \rangle$.*

Proof. Likewise in the proof of the necessary condition of Proposition 2, since we only used that $\ker S$ has a basis as (3). $\square$

This condition is not enough for S to be a glued semigroup, because the generating set $B_1 \cup B_2$ could be non-minimal. For example, if one get the numerical semigroups $T_1 = \langle 3, 5 \rangle$, $T_2 = \langle 2, 7 \rangle$ and $(\gamma_X, \gamma_Y) = (1, 0, 2, 0)$, one have the matrix as (5)

$$\left(\begin{array}{cc|cc} 5 & -3 & 0 & 0 \\ \hline 0 & 0 & 7 & -2 \\ \hline 1 & 0 & -2 & 0 \end{array} \right)$$

and the set $B_1 \cup B_2$ is $\{12, 20, 6, 21\}$ that it is not a minimal generating set.

Next corollary is devoted to solve this issue.

Corollary 10. *S is a glued semigroup if*

$$\left(\sum_{i=1}^{r} \gamma_{Xi} \right) \left(\sum_{i=1}^{t} \gamma_{Yi} \right) > 1. \tag{6}$$

Proof. Suppose that the set of generators $B_1 \cup B_2$ of S is non-minimal. Then one of its elements is a natural combination of the rest. Without loss of generality, we assume that this is the first element of $B_1 \cup B_2$.

Then there exist $\lambda_2, \ldots, \lambda_{r+t} \in \mathbb{N}$ such that

$$B_1(1, -\lambda_2, \ldots, -\lambda_r) = B_2(\lambda_{r+1}, \ldots, \lambda_{r+t}) \in G(\langle B_1 \rangle) \cap G(\langle B_2 \rangle).$$

By Proposition 9, there exists $\lambda \in \mathbb{Z}$ satisfying the equalities $B_1(1, -\lambda_2, \ldots, -\lambda_r) = B_2(\lambda_{r+1}, \ldots, \lambda_{r+t}) = B_1(\lambda\gamma_X)$. Also $\lambda \geq 0$ because $B_2(\lambda_{r+1}, \ldots, \lambda_{r+t}) \in S$, and consequently we obtain that

$$\nu = (1 - \lambda\gamma_{X1}, \underbrace{-\lambda_2 - \lambda\gamma_{X2}, \ldots, -\lambda_r - \lambda\gamma_{Xr}}_{\leq 0}) \in \ker(\langle B_1 \rangle) = \ker T_1.$$

---

[2] Note that $\gamma_X \notin \ker T_1$ and $\gamma_Y \notin \ker T_2$ because these semigroups are reduced.

- If $\lambda\gamma_{X1} = 0$, T_1 is not minimally generated, but it is not possible by hypothesis.
- If $\lambda\gamma_{X1} > 1$, then $0 > \nu \in \ker T_1$, but this is not possible because T_1 is a reduced semigroup.
- If $\lambda\gamma_{X1} = 1$, then $\lambda = \gamma_{X1} = 1$ and

  $$\nu = (0, \underbrace{-\lambda_2 - \gamma_{X2}, \ldots, -\lambda_r - \gamma_{Xr}}_{\leq 0}) \in \ker T_1.$$

  Again, if $\lambda_i + \gamma_{Xi} \neq 0$ for some $i = 2, \ldots, r$, T_1 is not a reduced semigroup. Then $\lambda_i = \gamma_{Xi} = 0$ for all $i = 2, \ldots, r$.

We have just proved that $\gamma_X = (1, 0, \ldots, 0)$.

In the general case, if S is not minimally generated it is because either γ_X or γ_Y are elements in the canonical bases of $\mathbb{N}^r$ or $\mathbb{N}^t$, respectively. To avoid this situation, it is sufficient to take γ_X and γ_Y satisfying $\left(\sum_{i=1}^{r} \gamma_{Xi}\right)\left(\sum_{i=1}^{t} \gamma_{Yi}\right) > 1$. □

From the above result we obtain a characterization of the glued semigroups: S is a glued semigroup if and only if $\ker S$ has a basis as (3) satisfying the condition (6).

Example 11. *Let* $T_1 = \langle(-7,2),(11,1),(5,0),(0,1)\rangle \subset \mathbb{Z}^2$ *and* $T_2 = \langle 3,5,7\rangle \subset \mathbb{N}$ *be two reduced affine semigroups.*

We compute their associated lattices

$$\ker T_1 = \langle(1,2,-3,-4),(2,-1,5,-3)\rangle \text{ and } \ker T_2 = \langle(-4,1,1),(-7,0,3)\rangle.$$

If we take $\gamma_X = (2,0,2,0)$ *and* $\gamma_Y = (1,2,1)$, *and construct the matrix*

$$\begin{pmatrix} 1 & 2 & -3 & -4 & 0 & 0 & 0 \\ 2 & -1 & 5 & -3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -4 & 1 & 1 \\ 0 & 0 & 0 & 0 & -7 & 0 & 3 \\ 2 & 0 & 2 & 0 & -1 & -2 & -1 \end{pmatrix},$$

we have the semigroup, $S \subset \mathbb{Z}_4 \times \mathbb{Z}^2$, *generated by*

$$\{\underbrace{(9,-5,35),(-17,12,-55),(-7,5,-25),(0,1,0)}_{B_1},\underbrace{(2,0,3),(2,0,5),(2,0,7)}_{B_2}\}.$$

This semigroup verifies that $\ker S$ *is generated by the rows of the above matrix, and it is the gluing of the semigroups* $\langle B_1\rangle$ *and* $\langle B_2\rangle$. *The ideal* $I_S \subset \mathbb{C}[x_1,\ldots,x_4,y_1,\ldots,y_3]$ *is generated*[3] *by*

$$\{x_1x_3^8x_4 - x_2^3,\ x_1x_2^2 - x_3^3x_4^4,\ x_1^2x_3^5 - x_2x_4^3, x_1^3x_2x_3^2 - x_7^7,$$
$$y_1y_3 - y_2^2,\ y_1^3y_2 - y_3^2,\ y_1^4 - y_2y_3,\ \underbrace{x_1^2x_3^2 - y_1^5y_2}_{\textit{glued binomial}}\},$$

then S *is really a glued semigroup.*

---

[3] See [14] to compute I_S when S has torsion.

In this way, we provide a procedure that allows us the construction of (glued) semigroups that are complete intersections. Regarding the following Lemma, it is sufficient that the semigroups T_1 and T_2 are complete intersections in order to be S as well. Next we give an algorithm to generate many examples of complete intersection semigroups.

Lemma 12. *T_1 and T_2 are two complete intersection semigroups if and only if S is complete intersection semigroup.*

Proof. It is trivial, since $\operatorname{rank}(\ker S) = \operatorname{rank}(\ker T_1) + \operatorname{rank}(\ker T_2) + 1$. $\square$

3.1 Generating affine glued semigroups

As one can check in Example 11, the semigroup S is not necessarily torsion-free. In general, a semigroup T is affine, i.e. it is torsion-free, if and only if the *invariant factors*[4] of the matrix whose rows are a basis of $\ker T$ are equal to one. We suppose that if the Smith Normal Form, D, of a matrix has some zero-columns, which are on the right side of D.

We use this fact to give conditions for S being torsion-free.

Let P_1, P_2, Q_1 and Q_2 some matrices with determinant ± 1 (i.e. unimodular matrices) such that $D_1 = P_1 L_1 Q_1$ and $D_2 = P_2 L_2 Q_2$ are the Smith Normal Form of L_1 and L_2, respectively.

If T_1 and T_2 are two affine semigroups, the invariant factors of L_1 and L_2 are equal to 1. Then

$$\left(\begin{array}{c|c} D_1 & 0 \\ \hline 0 & D_2 \\ \hline \gamma'_X & \gamma'_Y \end{array}\right) = \left(\begin{array}{c|c|c} P_1 & 0 & 0 \\ \hline 0 & P_2 & 0 \\ \hline 0 & 0 & 1 \end{array}\right) \underbrace{\left(\begin{array}{c|c} L_1 & 0 \\ \hline 0 & L_2 \\ \hline \gamma_X & -\gamma_Y \end{array}\right)}_{=:A} \left(\begin{array}{c|c} Q_1 & 0 \\ \hline 0 & Q_2 \end{array}\right), \tag{7}$$

where $\gamma'_X = \gamma_X Q_1$ and $\gamma'_Y = -\gamma_Y Q_2$. Let s_1 and s_2 be the numbers of zero-columns of D_1 and D_2 ($s_1, s_2 > 0$ because T_1 and T_2 are reduced).

Lemma 13. *S is an affine semigroup if and only if*

$$\gcd\left(\{\gamma'_{Xi}\}_{i=r-s_1}^{r} \cup \{\gamma'_{Yi}\}_{i=t-s_2}^{t}\right) = 1.$$

Proof. With the conditions imposed to T_1, T_2 and (γ_X, γ_Y), $\gcd\left(\{\gamma'_{Xi}\}_{i=r-s_1}^{r} \cup \{\gamma'_{Yi}\}_{i=t-s_2}^{t}\right) = 1$ is a necessary and sufficient condition for the invariant factors of A to be all equal to one. $\square$

In the following Corollary we give the explicit conditions that γ_X and γ_Y must fulfill to construct an affine semigroup.

Corollary 14. *S is an affine glued semigroup if and only if:*

1. *T_1 and T_2 are two affine semigroups.*
2. *$(\gamma_X, \gamma_Y) \in \mathbb{N}^{r+t}$.*

---

[4] The invariant factors of a matrix are the diagonal elements of its Smith Normal Form (see [2, Chapter 2] and [11, Chapter 2]).

3. $\sum_{i=1}^{r}\gamma_{Xi}$, $\sum_{i=1}^{t}\gamma_{Yi} > 1$.

4. There exist $f_{r-s_1},\dots,f_r,g_{t-s_2},\dots,g_t \in \mathbb{Z}$ *such that*

$$(f_{r-s_1},\dots,f_r)\cdot(\gamma'_{X(r-s_1)},\dots,\gamma'_{Xr})+(g_{t-s_2},\dots,g_t)\cdot(\gamma'_{Y(t-s_2)},\dots,\gamma'_{Yt}) = 1.$$

Proof. Trivial by construction, Corollary 10 and Lemma 13. ☐

Therefore, to construct an affine glued semigroup, it is enough to take two affine semigroups, and any solution, (γ_X, γ_Y), of the equations of the above corollary.

Example 15. *Let T_1 and T_2 be the semigroups of Example 11.*

We compute two elements $\gamma_X = (a_1, a_2, a_3, a_4)$ and $\gamma_Y = (b_1, b_2, b_3)$ in order to obtain an affine semigroup.

First of all, we perform a decomposition of the matrix as (7) by computing the integer Smith Normal Form of L_1 and L_2 :

$$\left(\begin{array}{cccc|ccc}
1 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & 0 & 0\\
\hline
0 & 0 & 0 & 0 & 1 & 0 & 0\\
0 & 0 & 0 & 0 & 0 & 1 & 0\\
\hline
a_1 & a_1-2a_2-a_3 & -7a_1+11a_2+5a_3 & 2a_1+a_2+a_4 & -b_1 & b_1+2b_2+3b_3 & -3b_1-5b_2-7b_3
\end{array}\right) =$$

$$\left(\begin{array}{cc|cc|c}
1 & 0 & 0 & 0 & 0\\
2 & -1 & 0 & 0 & 0\\
\hline
0 & 0 & -2 & 1 & 0\\
0 & 0 & 7 & -4 & 0\\
\hline
0 & 0 & 0 & 0 & 1
\end{array}\right)
\left(\begin{array}{cccc|ccc}
1 & 2 & -3 & -4 & 0 & 0 & 0\\
2 & -1 & 5 & -3 & 0 & 0 & 0\\
\hline
0 & 0 & 0 & 0 & -4 & 1 & 1\\
0 & 0 & 0 & 0 & -7 & 0 & 3\\
\hline
a_1 & a_2 & a_3 & a_4 & -b_1 & -b_2 & -b_3
\end{array}\right)
\left(\begin{array}{cccc|ccc}
1 & 1 & -7 & 2 & 0 & 0 & 0\\
0 & -2 & 11 & 1 & 0 & 0 & 0\\
0 & -1 & 5 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 1 & 0 & 0 & 0\\
\hline
0 & 0 & 0 & 0 & 1 & -1 & 3\\
0 & 0 & 0 & 0 & 0 & -2 & 5\\
0 & 0 & 0 & 0 & 0 & -3 & 7
\end{array}\right)$$

Second, by Corollary 14, we must find a solution to the system:

$$\left\{\begin{array}{l}
a_1+a_2+a_3+a_4 > 1\\
b_1+b_2+b_3 > 1\\
f_1, f_2, g_1 \in \mathbb{Z}\\
f_1(-7a_1+11a_2+5a_3)+f_2(2a_1+a_2+a_4)+g_1(-3b_1-5b_2-7b_3) = 1
\end{array}\right. ,$$

with $a_1, a_2, a_3, a_4, b_1, b_2, b_3 \in \mathbb{N}$. We compute a solution (in less than a second) using `FindInstance` *of Wolfram Mathematica (see [15]):*

$$\texttt{FindInstance}[(-7a_1{+}11a_2{+}5a_3){*}f_1{+}(2a_1{+}a_2{+}a_4){*}f_2{+}(3b_1{-}5b_2{-}7b_3){*}g_1 == 1$$

$$\&\&\, a1{+}a2{+}a3{+}a4 > 1\&\&\, b_1{+}b_2{+}b_3 > 1\&\&\, a_1 \geq 0\&\&\, a_2 \geq 0\&\&\, a_3 \geq 0\&\&\, a_4 \geq 0$$

$$\&\&\, b_1 \geq 0\&\&\, b_2 \geq 0\&\&\, b_3 \geq 0, \{a_1, a_2, a_3, a_4, b_1, b_2, b_3, f_1, f_2, g_1\},\ \mathit{Integers}\ ]$$

$$\Downarrow$$

$$\{\{a_1 \to 10, a_2 \to 1, a_3 \to 12, a_4 \to 0, b_1 \to 14, b_2 \to 0, b_3 \to 6, f_1 \to 1, f_2 \to 0, g_1 \to 0\}\}$$

Now, we take $\gamma_X = (10, 1, 12, 0)$ and $\gamma_Y = (14, 0, 6)$, and construct the matrix

$$A = \left(\begin{array}{ccccccc}
1 & 2 & -3 & -4 & 0 & 0 & 0\\
2 & -1 & 5 & -3 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & -4 & 1 & 1\\
0 & 0 & 0 & 0 & -7 & 0 & 3\\
10 & 1 & 12 & 0 & -14 & 0 & -6
\end{array}\right).$$

We have that $S \subset \mathbb{Z}^2$ *the affine semigroup, minimally generated by*

$$\{\underbrace{(149,-588),(-230,924),(-105,420),(1,0)}_{B_1},\underbrace{(0,3),(0,5),(0,7)}_{B_2}\},$$

satisfies that $\ker S$ *is generated by the rows of* A, *and it is the result of gluing the semigroups* $\langle B_1\rangle$ *and* $\langle B_2\rangle$. *The ideal* I_S *is generated by*

$$\{x_1x_3^8x_4 - x_2^3,\, x_1x_2^2 - x_3^3x_4^4,\, x_1^2x_3^5 - x_2x_4^3, x_1^3x_2x_3^2 - x_7^7,$$

$$y_1y_3 - y_2^2,\, y_1^3y_2 - y_3^2,\, y_1^4 - y_2y_3,\, \underbrace{x_1^{10}x_2x_3^{12} - y_1^{28}}_{\textit{glued binomial}}\},$$

therefore, S is a glued semigroup.

All the glued semigroups have been computed by using our programm `ecuaciones` which is available in [3] (this programm requires Wolfram Mathematica to run).

References

[1] E. Briales, A. Campillo, C. Marijuán, P. Pisón. *Minimal Systems of Generetors for Ideals of Semigroups.* J. Pure Appl. Algebra **124** (1998), 7–30.

[2] H. Cohen. *A Course in Computational Algebraic Number Theory,* Graduate Texts in Mathematics, **138**, Springer-Verlag, 1996.

[3] `ecuaciones`. `http://www.uca.es/dpto/C101/pags-personales/alberto.vigneron/1ecuaciones.rar`

[4] S. Eliahou. *Courbes monomiales et algébre de Rees symbolique.* PhD Thesis. Université of Genève, 1983.

[5] P. A. Garcia-Sanchez, I. Ojeda. *Uniquely presented finitely generated commutative monoids.* Pacific J. Math. **248**(1) (2010), 91–105.

[6] J. Herzog. *Generators and relations of abelian semigroups and semigroup rings.* Manuscripta Math. **3** (1970) 175–193.

[7] I. Ojeda, A. Vigneron-Tenorio. *Simplicial complexes and minimal free resolution of monomial algebras.* J. Pure Appl. Algebra **214** (2010), 850–861.

[8] I. Ojeda, A. Vigneron-Tenorio. *Indispensable binomials in semigroup ideals.* Proc. Amer. Math. Soc. **138** (2010), 4205–4216.

[9] H. Ohsugi, T. Hibi. *Toric ideals arising from contingency tables.* Commutative Algebra and Combinatorics. Ramanujan Mathematical Society Lecture Notes Series, Vol. 4, Ramanujan Mathematical Society, Mysore, India, 2007, pp. 91-115.

[10] J.C. Rosales. *On presentations of subsemigroups of* $\mathbb{N}^n$. Semigroup Forum **55** (1997), no. 2, 152–159.

[11] J.C. Rosales, P.A. García-Sánchez. *Finitely generated commutative monoids.* Nova Science Publishers, Inc., New York, 1999.

[12] B. Sturmfels. *Gröbner bases and convex polytopes*, volume 8 of *University Lecture Series.* American Mathematical Society, Providence, RI, 1996.

[13] A. Thoma. *Construction of Set Theoretic Complete Intersections via Semigroup Gluing.* Beiträge Algebra Geom. **41**(1) (2000), 195–198.

[14] A. Vigneron-Tenorio. *Semigroup Ideals and Linear Diophantine Equations.* Linear Algebra and its Applications **295** (1999), 133–144.

[15] `Wolfram Mathematica. http://www.wolfram.com/mathematica/`